

Cosmological Constant in SUGRA Models Inspired by Degenerate Vacua

C. D. Froggatt, R. Nevzorov

University of Glasgow, Glasgow, G12 8QQ, UK

H. B. Nielsen

The Niels Bohr Institute, Copenhagen, DK 2100, Denmark

It is well known that global symmetries protect local supersymmetry and a zero value for the cosmological constant in no-scale supergravity. A particular breakdown of these symmetries, which ensures the vanishing of the vacuum energy density, leads to the natural realisation of the multiple point principle (MPP). In the MPP inspired SUGRA models the cosmological constant is naturally tiny.

1. INTRODUCTION

Present astrophysical and cosmological observations indicate that there is a tiny energy density (the cosmological constant) $\Lambda \sim 10^{-123} M_{Pl}^4 \sim 10^{-55} M_Z^4$ spread all over the Universe, which is responsible for its accelerated expansion. This energy density is incredibly small as compared with the contribution of QCD condensates to the total vacuum energy density, which is expected to be of order $\Lambda_{QCD}^4 \simeq 10^{-74} M_{Pl}^4$. An even much larger contribution $\sim v^4 \simeq 10^{-62} M_{Pl}^4$ must come from the electroweak symmetry breaking, while the contribution of zero-modes should push the total vacuum energy density up to $\sim M_{Pl}^4$. Because of the enormous cancellation between the contributions of different condensates to Λ , the smallness of the cosmological constant should be regarded as a fine-tuning problem.

The smallness of the cosmological constant could be related to an almost exact symmetry. An exact global supersymmetry (SUSY) ensures zero value for the energy density at the minimum of the potential of the scalar fields. However the breakdown of supersymmetry induces a huge and positive contribution to the total vacuum energy density of order M_S^4 , where M_S is the SUSY breaking scale. The non-observation of superpartners of quarks and leptons implies that $M_S \gg 100$ GeV.

2. NO-SCALE SUPERGRAVITY AND THE MULTIPLE POINT PRINCIPLE

In ($N = 1$) supergravity (SUGRA) models the scalar potential is specified in terms of the Kähler function

$$G(\phi_M, \phi_M^*) = K(\phi_M, \phi_M^*) + \ln |W(\phi_M)|^2, \quad (1)$$

which is a combination of two functions: the Kähler potential $K(\phi_M, \phi_M^*)$ and the superpotential $W(\phi_M)$. Here we use standard supergravity mass units: $\frac{M_{Pl}}{\sqrt{8\pi}} = 1$. The SUGRA scalar potential is given by

$$V(\phi_M, \phi_M^*) = \sum_{M, \bar{N}} e^G \left(G_M G^{\bar{M}\bar{N}} G_{\bar{N}} - 3 \right) + \frac{1}{2} \sum_a (D^a)^2, \quad D^a = g_a \sum_{i,j} (G_i T_{ij}^a \phi_j), \quad (2)$$

$$G_M \equiv \partial G / \partial \phi_M, \quad G_{\bar{M}} \equiv \partial G / \partial \phi_M^*, \quad G_{\bar{N}M} \equiv \frac{\partial^2 G}{\partial \phi_M^* \partial \phi_M}, \quad G^{M\bar{N}} = G_{\bar{N}M}^{-1},$$

where g_a is the gauge coupling constant associated with the generator T^a of the gauge transformations. In order to break supersymmetry in ($N = 1$) SUGRA models, a hidden sector is introduced. It is assumed that the superfields of the hidden sector (z_i) interact with the observable ones only by means of gravity. At the minimum of the scalar potential, hidden sector fields acquire vacuum expectation values breaking local SUSY and generating a non-zero gravitino mass $m_{3/2} = \langle e^{G/2} \rangle$.

In general the vacuum energy density at the minimum of the SUGRA scalar potential (2) is huge and negative $\rho_\Lambda \sim -m_{3/2}^2 M_{Pl}^2$, so that an enormous fine-tuning is required to keep ρ_Λ around the observed value of the cosmological constant. The situation changes dramatically if supergravity is supplemented by a global symmetry that ensures a zero value for the cosmological constant. This is precisely what happens in no-scale supergravity. The Lagrangian

of a no-scale SUGRA model is invariant under the imaginary translations and dilatations which are subgroups of the $SU(1,1)$ group. The simplest no-scale SUGRA model involves only one hidden sector superfield T and a set of chiral supermultiplets φ_σ in the observable sector, which transform differently under the imaginary translations ($T \rightarrow T + i\beta$, $\varphi_\sigma \rightarrow \varphi_\sigma$) and dilatations ($T \rightarrow \alpha^2 T$, $\varphi_\sigma \rightarrow \alpha \varphi_\sigma$). These symmetries constrain the Kähler function so that

$$K = -3 \ln \left[T + \bar{T} - \sum_\sigma \zeta_\sigma |\varphi_\sigma|^2 \right], \quad W = \sum_{\sigma, \lambda, \gamma} \frac{1}{6} Y_{\sigma\lambda\gamma} \varphi_\sigma \varphi_\lambda \varphi_\gamma. \quad (3)$$

The structure of the Kähler function (3) ensures the vanishing of the vacuum energy density in the considered model. Thus imaginary translations and dilatations protect a zero value for the cosmological constant in supergravity. However these symmetries also preserve supersymmetry in all vacua, which has to be broken in any phenomenologically acceptable theory.

It was argued that the breakdown of dilatation invariance does not necessarily result in a non-zero vacuum energy density [1]–[2]. Let us consider a SUGRA model with two hidden sector fields T and z . Suppose that T and the observable superfields φ_σ transform under imaginary translations and dilatations as before, whereas the superfield z transforms similarly to φ_σ . Then the hidden sector superfield z can appear in the full superpotential of the model:

$$W(z, \varphi_\alpha) = \kappa \left(z^3 + \mu_0 z^2 + \sum_{n=4}^{\infty} c_n z^n \right) + \sum_{\sigma, \beta, \gamma} \frac{1}{6} Y_{\sigma\beta\gamma} \varphi_\sigma \varphi_\beta \varphi_\gamma. \quad (4)$$

Here we include a bilinear mass term for the superfield z and higher order terms $c_n z^n$ that spoil dilatation invariance. At the same time we do not allow the breakdown of dilatation invariance in the superpotential of the observable sector, in order to avoid the appearance of potentially dangerous terms which lead, for instance, to the so-called μ -problem in the simplest SUSY models.

We also assume that the dilatation invariance is broken in the Kähler potential of the observable sector, so that the full Kähler potential takes the form:

$$K = -3 \ln \left[T + \bar{T} - |z|^2 - \sum_\sigma \zeta_\sigma |\varphi_\sigma|^2 \right] + \sum_{\sigma, \lambda} \left(\frac{\eta_{\sigma\lambda}}{2} \varphi_\sigma \varphi_\lambda + h.c. \right) + \sum_\sigma \xi_\sigma |\varphi_\sigma|^2. \quad (5)$$

Here we restrict our consideration to the simplest set of terms that break dilatation invariance in the Kähler potential. Additional terms which are proportional to $|\varphi_\alpha|^2$ normally appear in minimal SUGRA models. The other terms $\eta_{\alpha\beta} \varphi_\alpha \varphi_\beta$ give rise to effective μ terms after the spontaneous breakdown of local supersymmetry, solving the μ problem. We only allow the breakdown of the dilatation invariance in the Kähler potential of the observable sector, since any variations in the Kähler potential of the hidden sector may spoil the vanishing of the vacuum energy density in global minima.

In the considered SUGRA model the scalar potential of the hidden sector is positive definite

$$V(T, z) = \frac{1}{3(T + \bar{T} - |z|^2)^2} \left| \frac{\partial W(z)}{\partial z} \right|^2, \quad (6)$$

so that the vacuum energy density vanishes near its global minima. The minima of the scalar potential (6) are attained at the stationary points of the hidden sector superpotential. In the simplest case when all the coefficients $c_n = 0$, $W(z)$ has two extremum points at $z = 0$ and $z = -\frac{2\mu_0}{3}$. In the first vacuum, where $z = -\frac{2\mu_0}{3}$, local supersymmetry is broken so that the gravitino becomes massive

$$m_{3/2} = \left\langle \frac{W(z)}{(T + \bar{T} - |z|^2)^{3/2}} \right\rangle = \frac{4\kappa\mu_0^3}{27 \left\langle \left(T + \bar{T} - \frac{4\mu_0^2}{9} \right)^{3/2} \right\rangle}. \quad (7)$$

and all scalar particles get non-zero masses $m_\sigma \sim \frac{m_{3/2} \xi_\sigma}{\zeta_\sigma}$. In the second minimum, with $z = 0$, the superpotential of the hidden sector vanishes and local SUSY remains intact, so that the low-energy limit of this theory is described

by a pure SUSY model in flat Minkowski space. If the high order terms $c_n z^n$ are present in Eq. (4), the scalar potential of the hidden sector may have many degenerate vacua with broken and unbroken supersymmetry in which the vacuum energy density vanishes.

Thus the considered breakdown of dilatation invariance leads to a natural realisation of the multiple point principle (MPP). The MPP postulates the existence of many phases with the same energy density which are allowed by a given theory [3]. When applied to ($N = 1$) supergravity, MPP implies the existence of a vacuum in which the low-energy limit of the considered theory is described by a pure supersymmetric model in flat Minkowski space. According to the MPP this vacuum and the physical one in which we live must be degenerate. Such a second vacuum is only realised if the SUGRA scalar potential has a minimum where the following conditions are satisfied

$$\left\langle W(z) \right\rangle = \left\langle \frac{\partial W(z)}{\partial z} \right\rangle = 0. \quad (8)$$

This would normally require an extra fine-tuning. However, in the SUGRA models considered above, the MPP conditions are fulfilled automatically without any extra fine-tuning at the tree-level.

3. COSMOLOGICAL CONSTANT IN THE MPP INSPIRED SUGRA MODELS

Since the vacuum energy density of supersymmetric states in flat Minkowski space is just zero and all vacua in the MPP inspired SUGRA models are degenerate, the cosmological constant problem is thereby solved to first approximation by assumption. However the value of the cosmological constant may differ from zero in the considered models. This occurs if non-perturbative effects in the observable sector give rise to the breakdown of supersymmetry in the second vacuum (phase). Our MPP philosophy then requires that the physical phase in which local supersymmetry is broken in the hidden sector has the same energy density as a phase where supersymmetry breakdown takes place in the observable sector.

If supersymmetry breaking takes place in the second vacuum, it is caused by the strong interactions. When the gauge couplings at high energies are identical in both vacua, their running down to the scale $M_S \simeq m_{3/2}$ are also the same. Below the scale M_S all superparticles in the physical vacuum decouple and the corresponding beta functions change. Using the value of $\alpha_3^{(1)}(M_Z) \approx 0.118 \pm 0.003$ and the matching condition $\alpha_3^{(2)}(M_S) = \alpha_3^{(1)}(M_S)$, one finds the scale Λ_{SQCD} , where the supersymmetric QCD interactions become strong in the second vacuum:

$$\Lambda_{SQCD} = M_S \exp \left[\frac{2\pi}{b_3 \alpha_3^{(2)}(M_S)} \right], \quad \frac{1}{\alpha_3^{(2)}(M_S)} = \frac{1}{\alpha_3^{(1)}(M_Z)} - \frac{\tilde{b}_3}{4\pi} \ln \frac{M_S^2}{M_Z^2}. \quad (9)$$

In Eq.(9) $\alpha_3^{(1)}$ and $\alpha_3^{(2)}$ are the values of the strong gauge couplings in the physical and second minima of the SUGRA scalar potential, while $\tilde{b}_3 = -7$ and $b_3 = -3$ are the one-loop beta functions of the SM and MSSM. At the scale Λ_{SQCD} the t-quark Yukawa coupling in the MSSM is of the same order of magnitude as the strong gauge coupling. The large Yukawa coupling of the top quark may result in the formation of a quark condensate that breaks supersymmetry inducing a non-zero positive value for the cosmological constant $\Lambda \simeq \Lambda_{SQCD}^4$.

In Fig. 1 the dependence of Λ_{SQCD} on the SUSY breaking scale M_S is examined. Because $\tilde{b}_3 < b_3$ the QCD gauge coupling below M_S is larger in the physical minimum than in the second one. Therefore the value of Λ_{SQCD} is much lower than the QCD scale in the Standard Model and diminishes with increasing M_S . When the supersymmetry breaking scale in our vacuum is of the order of 1 TeV, we obtain $\Lambda_{SQCD} = 10^{-26} M_{Pl} \simeq 100$ eV. This results in an enormous suppression of the total vacuum energy density ($\Lambda \simeq 10^{-104} M_{Pl}^4$) compared to say an electroweak scale contribution in our vacuum $v^4 \simeq 10^{-62} M_{Pl}^4$. From the rough estimate $\Lambda \simeq \Lambda_{SQCD}^4$ of the energy density, it can be easily seen that the measured value of the cosmological constant is reproduced when $\Lambda_{SQCD} = 10^{-31} M_{Pl} \simeq 10^{-3}$ eV [1], [4]. The appropriate values of Λ_{SQCD} can therefore only be obtained for $M_S = 10^3 - 10^4$ TeV. However the consequent large splitting within SUSY multiplets would spoil gauge coupling unification in the MSSM and reintroduce the hierarchy problem, which would make the stabilisation of the electroweak scale rather problematic.




Figure 1: The value of $\log[\Lambda_{SQCD}/M_{Pl}]$ versus $\log M_S$. The thin and thick solid lines correspond to the pure MSSM and the MSSM with an extra pair of $\mathbf{5} + \bar{\mathbf{5}}$ multiplets. The dashed and dash-dotted lines represent the uncertainty in $\alpha_3(M_Z)$, i.e. $\alpha_3(M_Z) = 0.112 - 0.124$. The horizontal line corresponds to the observed value of $\Lambda^{1/4}$. The SUSY breaking scale M_S is given in GeV.

A model consistent with electroweak symmetry breaking and cosmological observations can be constructed, if the MSSM particle content is supplemented by an additional pair of $\mathbf{5} + \bar{\mathbf{5}}$ multiplets. These new bosons and fermions would not affect gauge coupling unification, because they form complete representations of $SU(5)$. In the physical vacuum these extra particles would gain masses around the supersymmetry breaking scale. The corresponding mass terms in the superpotential are generated after the spontaneous breaking of local supersymmetry, due to the presence of the bilinear terms $[\eta(\mathbf{5} \cdot \bar{\mathbf{5}}) + h.c.]$ in the Kähler potential of the observable sector. Near the second minimum of the SUGRA scalar potential the new particles would be massless, since $m_{3/2} = 0$. Therefore they give a considerable contribution to the β functions ($b_3 = -2$), reducing Λ_{SQCD} further. In this case the observed value of the cosmological constant can be reproduced even for $M_S \simeq 1$ TeV (see Fig. 1) [1], [4].

Acknowledgments

We would like to thank A. Faraggi, D.R.T. Jones, S.F. King, A. Pilaftsis and J. Polchinski for fruitful discussions. RN acknowledges support from the SHEFC grant HR03020 SUPA 36878.

References

- [1] C.D. Froggatt, R. Nevzorov, H.B. Nielsen, “On the smallness of the cosmological constant in SUGRA models”, Nucl. Phys. B **743** (2006) 133; C.D. Froggatt, R. Nevzorov, H.B. Nielsen, “Smallness of the cosmological constant and the multiple point principle”, J. Phys. Conf. Ser. **110** (2008) 072012.
- [2] C.D. Froggatt, L. Laperashvili, R. Nevzorov, H.B. Nielsen, “No-scale supergravity and the multiple point principle”, arXiv:hep-ph/0411273.
- [3] D.L. Bennett, H.B. Nielsen, “Predictions for nonAbelian finestructure constants from multicriticality”, Int. J. Mod. Phys. A **9** (1994) 5155; D.L. Bennett, C.D. Froggatt, H.B. Nielsen, “Nonlocality as an explanation for fine tuning and field replication in nature”, arXiv:hep-ph/9504294.
- [4] C.D. Froggatt, L. Laperashvili, R. Nevzorov, H.B. Nielsen, “Cosmological constant in SUGRA models and the multiple point principle”, Phys. Atom. Nucl. **67** (2004) 582.